

GEOMETRY OF QUIVER GRASSMANNIANS OF KRONECKER TYPE AND CANONICAL BASIS OF CLUSTER ALGEBRAS

GIOVANNI CERULLI IRELLI AND FRANCESCO ESPOSITO

ABSTRACT. We study quiver Grassmannians associated with indecomposable representations of the Kronecker quiver. We find a cellular decomposition of them and we compute their Betti numbers. As an application, we find a geometric realization of the “canonical basis” of cluster algebras of type $A_1^{(1)}$ found by Sherman and Zelevinsky in [24] and of type $A_2^{(1)}$ found in [18].

1. INTRODUCTION

Cluster algebras are commutative $\mathbb{Z}$ -subalgebras of the field of rational functions in a finite number of indeterminates which have been introduced and studied by S. Fomin and A. Zelevinsky in a series of papers [13], [14], [2] and [16]. To every quiver Q without loops and 2-cycles it is associated a coefficient-free cluster algebra $\mathcal{A}_Q$. In [4], [6], [5] and [11] the authors describe the cluster variables of $\mathcal{A}_Q$ via a map, called the Caldero–Chapoton map, between the representations of Q and the field of rational functions in n variables (we address the reader to the survey [20]). Such map is defined in terms of Euler–Poincaré characteristic of some complex projective varieties attached to every representation M of Q and called quiver Grassmannians. By definition the quiver Grassmannian $Gr_{\mathbf{e}}(M)$ consists of all sub-representations of M of dimension vector $\mathbf{e}$. These varieties are considered in several places, e.g. [6], [5], [7], [17, section 12.3], [22], and in this paper we try to add some more geometric information about them, at least in the case of the Kronecker quiver. In [8] authors compute the Euler–Poincaré characteristic of quiver Grassmannians associated with the Kronecker quiver and they conjecture the existence of a cellular decomposition which we find here. In [19] a torus action on some quiver Grassmannians has been found and this allows to produce a cellular decomposition of them in the case they are smooth.

In this paper we study quiver Grassmannians associated with the Kronecker quiver. A representation of the Kronecker quiver (in the sequel we say a Q -representation) is a quadruple $M = (M_1, M_2, m_a, m_b)$ where M_1 and M_2 are finite dimensional complex vector spaces and $m_a, m_b : M_1 \rightarrow M_2$ are two linear maps between them. Given two non-negative integers e_1 and e_2 , the variety $Gr_{(e_1, e_2)}(M)$ is defined as the set

$$\{(N_1, N_2) \in Gr_{e_1}(M_1) \times Gr_{e_2}(M_2) : m_a(N_1) \subset N_2, m_b(N_1) \subset N_2\}$$

where $Gr_e(V)$ denotes the Grassmannians of e -dimensional vector spaces in a vector space V . This is a projective variety which is in general not

¹Research supported by grant CPDA071244/07 of Padova University

smooth. When $M_1 = M_2 = \mathbf{C}^n$ and $m_a = Id$ is the identity and $m_b = J_n(0)$ is an indecomposable nilpotent Jordan block, the representation M is regular indecomposable and we denote it by R_n . The corresponding quiver Grassmannians $Gr_{(e_1, e_2)}(R_n)$ are the main subjects of this paper. We briefly write X for one of these quiver Grassmannians. We find that a one-dimensional torus T acts on X (section 2.2). We provide a stratification of X (see section 2.5)

$$(1) \quad X = X_0 \supseteq X_1 \supseteq \cdots \supseteq X_s$$

for $s = \min(e_1, n - e_2)$ into closed subvarieties $X_k \simeq Gr_{(e_1-k, e_2-k)}(R_{n-2k})$. Moreover X_{k+1} is the singular locus of X_k and the difference $X_k \setminus X_{k+1}$ is a smooth quasi-projective variety which is not complete. As a consequence of the stratification (1) we get that X is smooth if and only if $s = 0$, i.e. either $e_1 = 0$ or $e_2 = n$, in which cases the quiver Grassmannian specializes to an usual Grassmannian of vector subspaces.

In section 2.6 we prove that Białyński-Birula's theorem on cellular decomposition of smooth projective varieties applies to $X_k \setminus X_{k+1}$ and we can hence prove that X decomposes

$$X = \cup_{L \in X^T} X_L$$

into attracting sets of T -fixed points

$$X_L := \{N \in X \mid \lim_{\lambda \rightarrow 0} t_\lambda N = L\}$$

and these sets are affine spaces. In section 2.7 we describe the cell X_L : if L is indecomposable then X_L is the orbit of L by the action of a unipotent group; if $L = L' \oplus L''$ with L' of "lower weight than" L'' then the following formula holds:

$$\dim X_{L' \oplus L''} = \dim X_{L'} + \dim X_{L''} - \langle \mathbf{dim} L', \mathbf{dim} L'' \rangle$$

where $\langle \cdot, \cdot \rangle$ denotes the Euler form of the Kronecker quiver. As a consequence of this formula we are able to compute the Poincaré polynomials of every indecomposable representation of the Kronecker quiver: let P_n (resp. I_n) be the indecomposable preprojective of dimension $(n, n+1)$ (resp. $(n+1, n)$). We have (see section 2.8)

$$(2) \quad \begin{aligned} P_{Gr_e(R_n)}(t) &= P_{Gr_{(e_2-e_1)}(e_2)}(t) P_{Gr_{(e_2-e_1)}(n-e_1)}(t) \\ P_{Gr_e(P_n)}(t) &= P_{Gr_{e_1}(e_2-1)}(t) P_{Gr_{(e_2-e_1)}(n+1-e_1)}(t) \\ P_{Gr_e(I_n)}(t) &= P_{Gr_{e_1}(e_2+1)}(t) P_{Gr_{(e_2-e_1)}(n-e_1)}(t) \end{aligned}$$

where $Gr_t(s)$ denotes the Grassmannian of t -dimensional vector subspaces of an s -dimensional vector space.

We remark that formulas (2) yield another proof of results of Szanto [25]. Indeed $Gr_e(P_n)$ and $Gr_e(I_n)$ are smooth over any field and hence this follows from [3]. For $X = Gr_e(R_n)$, both X and all the X_k 's may be defined over $\mathbb{Z}$ and the results of theorem 2.1 continues to hold over $\mathbb{Z}$. In particular $X_k \setminus X_{k+1}$ is smooth over $\mathbb{Z}$ and by standard theorems in l -adic cohomology one has that the cohomology of the quiver grassmannian over the field F_q is the same as in characteristic zero and the Frobenius acts by a suitable power of q . Applying the Lefschetz fixed point formula, our formula for the Poincaré polynomials provide another proof of the results in [25]. It would

be interesting to know if the cellular decomposition continues to hold in positive characteristic.

In section 3 we consider the (coefficient-free) cluster algebra $\mathcal{A}$ of type $A_1^{(1)}$. The cluster variables of $\mathcal{A}$ are the rational functions x_m , $m \in \mathbb{Z}$, of the field $\mathcal{F} = \mathbb{Q}(x_1, x_2)$ recursively generated by the following relation:

$$x_k x_{k+2} = x_{k+1}^2 + 1.$$

In this case the Caldero–Chapoton map $M \mapsto CC(M)$ associates to a Q -representation M of dimension vectors (d_1, d_2) the following Laurent polynomial:

$$(3) \quad CC(M) := \frac{\sum_{\mathbf{e}} \chi(Gr_{\mathbf{e}}(M)) x_1^{2(d_2 - e_2)} x_2^{2e_1}}{x_1^{d_1} x_2^{d_2}}$$

In [6] it is proved that the map $M \mapsto CC(M)$ restricts to a bijection between the indecomposable rigid Q -representations M (i.e. $Ext^1(M, M) = 0$) and the cluster variables of $\mathcal{A}$ different from x_1 and x_2 . Moreover it has the following property: $CC(M \oplus N) = CC(M)CC(N)$, under which cluster monomials not divisible by x_1 or x_2 , i.e. monomials of the form $x_k^a x_{k+1}^b$ for $k \in \mathbb{Z} \setminus \{1, 2\}$ and $a, b \geq 0$, are in bijection with rigid Q -representations.

In [24] the authors introduce distinguished elements $\{z_n \mid n \geq 1\}$ of $\mathcal{F}$ recursively defined by:

$$(4) \quad \begin{aligned} z_0 &= 2 \\ z_1 &= x_0 x_3 - x_1 x_2 \\ z_{n+1} &= z_1 z_n - z_{n-1} \quad n \geq 1 \end{aligned}$$

and they prove that the set

$$\mathbf{B} := \{\text{cluster monomials}\} \cup \{z_n : n \geq 1\}$$

is a $\mathbb{Z}$ -basis of $\mathcal{A}$ such that positive linear combinations of its elements coincide with the set of all *positive* elements of $\mathcal{A}$ (i.e. elements which are positive Laurent polynomials in every cluster of $\mathcal{A}$). They call this basis the canonical basis of $\mathcal{A}$. We give a geometric realization of $\mathbf{B}$ by using the Caldero–Chapoton map: cluster monomials are image of rigid representations and quiver Grassmannians associated with rigid quiver representations are smooth [7]. Having this in mind in section 3 we prove the following: for every $n \geq 1$:

$$z_n = \frac{\sum_{\mathbf{e}} \chi(Gr_{\mathbf{e}}(R_n)^{sm}) x_1^{2(n - e_2)} x_2^{2e_1}}{x_1^n x_2^n}$$

where $Gr_{\mathbf{e}}(R_n)^{sm} := X_0 \setminus X_1$ denotes the smooth part of $Gr_{\mathbf{e}}(R_n)$.

A similar construction can be made in a cluster algebra $\mathcal{A}_2$ of type $A_2^{(1)}$. These cluster algebras are studied in [18] and some results are recalled in section 3.2: the canonical basis of $\mathcal{A}_2$ consists of cluster monomials together with elements $\{u_n w^k, u_n z^k : k \geq 0, n \geq 1\}$ where w and z are two cluster variables and the u_n are defined similarly to (4) as follows:

$$(5) \quad \begin{aligned} u_0 &= 2 \\ u_1 &= zw - 2 \\ u_{n+1} &= u_1 u_n - u_{n-1} \quad n \geq 1 \end{aligned}$$

We prove that u_n is obtained by evaluating the Caldero–Chapoton map at the *smooth part* of the regular indecomposable representation of a quiver of type $A_2^{(1)}$ of dimension $(1, 1, 1)$ which lies in an homogeneous tube.

2. GEOMETRIC STRUCTURE OF QUIVER GRASSMANNIANS

Let

$$Q := 1 \begin{array}{c} \xrightarrow{a} \\ \xrightarrow{b} \end{array} 2$$

be the Kronecker quiver. As usual, we denote a complex Q –representation $M = (M_1, M_2, m_a, m_b)$ as follows:

$$M = M_1 \begin{array}{c} \xrightarrow{m_a} \\ \xrightarrow{m_b} \end{array} M_2 .$$

A sub–representation N of M consists of vector subspaces N_1 and N_2 of M_1 and M_2 respectively such that $m_a(N_1) \subset N_2$ and $m_b(N_1) \subset N_2$. The vector $\mathbf{dim}(M) := (\dim M_1, \dim M_2)$ is called the dimension vector of M . A morphism $g : M \rightarrow M'$ from a Q –representation M to a Q –representation M' is a couple (g_1, g_2) of linear maps $g_1 : M_1 \rightarrow M'_1$ and $g_2 : M_2 \rightarrow M'_2$ such that $m'_a \circ g_1 = g_2 \circ m_a$ and $m'_b \circ g_1 = g_2 \circ m_b$. The set of Q –representations form a category which is an abelian Krull–Schmidt category via the natural notions of direct sums, kernel and cokernel (see e.g. [1]). The classification of Q –representations which are *indecomposable* (i.e. that are not direct sum of two non–trivial sub–representations) goes back to Kronecker [21]. The following is their complete list: there are the *indecomposable preprojectives* (for $n \geq 0$):

$$P_n = k^n \begin{array}{c} \xrightarrow{\varphi_1} \\ \xrightarrow{\varphi_2} \end{array} k^{n+1}$$

where $k = \mathbf{C}$ denotes the field of complex numbers, φ_1, φ_2 are the two immersions respectively in the vector subspace spanned by the first and the last basis vectors. There are the *indecomposable regulars*:

$$R_n(\lambda) = k^n \begin{array}{c} \xrightarrow{Id} \\ \xrightarrow{J_n(\lambda)} \end{array} k^n , \quad R_n(\infty) = k^n \begin{array}{c} \xrightarrow{J_n(0)} \\ \xrightarrow{Id} \end{array} k^n$$

where $\lambda \in k$ and $J_n(\lambda)$ denotes the n –th indecomposable Jordan block of eigenvalue λ and Id denotes the identity matrix. And finally there are the *indecomposable preinjectives* (for $n \geq 0$):

$$I_n = k^{n+1} \begin{array}{c} \xrightarrow{\varphi_2^t} \\ \xrightarrow{\varphi_1^t} \end{array} k^n$$

where φ_1^t and φ_2^t are the transpose of the matrices φ_1 and φ_2 defined above. For an indecomposable Q –representation M we denote by $\mathcal{B}_i = \{v_k^{(i)}\}$, $i = 1, 2$, the standard basis of M_i with respect to which M has the previous presentation. All the other Q –representations are direct sums of these ones. Direct sums of indecomposable preprojectives (resp. regulars, preinjectives) are called preprojective (resp. regular, preinjective) Q –representations.

Given non-negative integers e_1 and e_2 and a Q -representation M we consider the variety:

$$Gr_{\mathbf{e}}(M) := \{N \leq M \mid \mathbf{dim}(N) = (e_1, e_2)\}$$

called the quiver Grassmannian of dimension $\mathbf{e} = (e_1, e_2)$ of M . This is closed inside the product $Gr_{e_1}(M_1) \times Gr_{e_2}(M_2)$ of usual Grassmannians of vector subspaces and it is hence a complex projective variety. In [7] it is shown that the tangent space $T_N(Gr_{\mathbf{e}}(M))$ at a point N of $Gr_{\mathbf{e}}(M)$ equals:

$$(6) \quad T_N(Gr_{\mathbf{e}}(M)) = Hom(N, M/N).$$

Moreover the following inequalities hold for $Z := Gr_{\mathbf{e}}(M)$:

$$(7) \quad \langle \mathbf{e}, \mathbf{d} - \mathbf{e} \rangle \leq \dim Z \leq \dim T_N(Z) \leq \langle \mathbf{e}, \mathbf{d} - \mathbf{e} \rangle + \dim Ext^1(M, M)$$

where $\langle (a, b)^t, (c, d)^t \rangle := ac + bd - 2ad$ is the Euler form of Q and $\mathbf{d} := \mathbf{dim} M$ so that $\mathbf{dim}(M/N) = \mathbf{d} - \mathbf{e}$. In particular if M is *rigid*, i.e. $Ext^1(M, M) = 0$, then all the quiver Grassmannians associated with it are smooth (see also [11, proposition 3.5] for another such result) and they have dimension $\dim Gr_{\mathbf{e}}(M) = \langle \mathbf{e}, \mathbf{d} - \mathbf{e} \rangle$. It is known that the rigid Q -representations are the following:

$$P_n^{\oplus a} \oplus P_{n+1}^{\oplus b}, \quad I_n^{\oplus a} \oplus I_{n+1}^{\oplus b}$$

for all $n \geq 0$ and $a, b \geq 0$.

2.1. Action of a group on quiver Grassmannians. Let M be a Q -representation of dimension vector $\mathbf{d} = (d_1, d_2)$ we consider the group

$$\mathcal{N}(M) := \{(A, B) \in \prod_{i=1}^2 GL_{d_i}(M_i) \mid \exists \lambda \in \mathbf{C}^* : m_a A = B m_a, m_b A = \lambda B m_b\}.$$

Note that the automorphism group of M is a closed subgroup of $\mathcal{N}(M)$ (for $\lambda = 1$). The group $\mathcal{N}(M)$ acts on $Gr_{\mathbf{e}}(M)$ as follows $(A, B) \cdot (N_1, N_2) := (AN_1, BN_2)$.

Lemma 2.0.1. (1) For every $n \geq 0$, $\mathcal{N}(P_n) \simeq \mathbf{C}^* \times \mathbf{C}^*$;
(2) For every $n \geq 0$, $\mathcal{N}(I_n) \simeq \mathbf{C}^* \times \mathbf{C}^*$.

Proof. It follows easily from the definition that $\mathcal{N}(P_n)$ consists of diagonal matrices (A, B) of the form:

$$A = \text{diag}(a, a\lambda, \dots, a\lambda^{n-1})$$

$$B = \text{diag}(a, a\lambda, \dots, a\lambda^n)$$

for $a, \lambda \in \mathbf{C}^*$. Similarly for $\mathcal{N}(I_n)$. □

Proposition 2.0.1. Let M be a rigid Q -representation. Then every quiver Grassmannian $Gr_{\mathbf{e}}(M)$ associated with M has a cellular decomposition.

Proof. The variety $Gr_{\mathbf{e}}(M)$ is smooth and $\mathcal{N}(M) \supset \mathbf{C}^* =: T$. The torus T acts with finitely many fixed points. It hence follows by Bialinicky–Birula results [3] (see also [9, section 2.4]) that it has a cellular decomposition into attracting sets of its T -fixed points. □

In the rest of the paper we mainly concentrate on the quiver Grassmannians associated with indecomposable regular Q -representations.

2.2. The variety $X = Gr_e(R_n)$. From now on we will focus on quiver Grassmannians associated with indecomposable regular Q -representations. It is not hard to show that $Gr_e(R_n(\lambda)) = Gr_e(R_n(\mu))$ for every $\lambda, \mu \in k \cup \{\infty\}$ (see e.g. [19]) and we hence consider the variety $X := Gr_e(R_n(0))$. It follows from the definition that

$$X = \{N_1 \subset N_2 \subset k^n \mid J_n(0)N_1 \subset N_2, \dim N_i = e_i, i = 1, 2\}$$

and hence X is a closed subvariety of a partial flag variety. The group $\mathcal{N} := \mathcal{N}(R_n)$ is the following:

$$\mathcal{N} := \{A \in GL_n(\mathbf{C}) \mid AJA^{-1} = \lambda J, \text{ for some } \lambda \in \mathbf{C}^*\}$$

where $J := J_n(0)$ and it acts on X as $A \cdot (N_1, N_2) = (AN_1, AN_2)$.

Lemma 2.0.2. *The group $\mathcal{N}$ is the semi-direct product*

$$\mathcal{N} = \mathcal{U} \rtimes T \times Z$$

where $\mathcal{U}$ is the unipotent radical of $\mathcal{N}$ of unipotent triangular Toeplitz matrices given by:

$$\mathcal{U} := \{\mathbf{1}_n + \sum_{i=1}^{n-1} a_i J_n^i(0) : a_i \in \mathbf{C}\},$$

T is the one-dimensional torus $T = \{t_\lambda : \lambda \in \mathbf{C}^*\}$ where t_λ is the diagonal matrix

$$t_\lambda := \text{diag}(1, \lambda, \lambda^2, \dots, \lambda^{n-1})$$

and Z consists of central elements $\{a_0 \mathbf{1}_n \mid a_0 \in \mathbf{C}^*\}$.

Proof. Every element of $\mathcal{UTZ}$ belongs to $\mathcal{N}$. Viceversa let $A \in \mathcal{N}$. Then it is easy to see that the columns $\mathbf{a}_1, \dots, \mathbf{a}_n$ of A satisfy the relation $\mathbf{a}_{k+1} = J\mathbf{a}_k$ and hence $A \in \mathcal{UTZ}$. It is now easy to see that $\mathcal{U}$ is normal in $\mathcal{UTZ}$. $\square$

For example for $n = 5$, an element A of the group $\mathcal{U}$ and a element t_λ of T have the form:

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ a_1 & 1 & 0 & 0 & 0 \\ a_2 & a_1 & 1 & 0 & 0 \\ a_3 & a_2 & a_1 & 1 & 0 \\ a_4 & a_3 & a_2 & a_1 & 1 \end{pmatrix} \quad t_\lambda = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \lambda & 0 & 0 & 0 \\ 0 & 0 & \lambda^2 & 0 & 0 \\ 0 & 0 & 0 & \lambda^3 & 0 \\ 0 & 0 & 0 & 0 & \lambda^4 \end{pmatrix}$$

for $a_1, a_2, a_3, a_4 \in \mathbf{C}$ and $\lambda \in \mathbf{C}$. Clearly $L \in X^T$ if and only if L is a coordinate sub-representation of R_n , i.e. both L_1 and L_2 are coordinate subspaces of k^n . In the next section we will encode this information in a combinatorial tool which is called the coefficient quiver of R_n .

We conclude this section by pointing out a useful isomorphism:

$$(8) \quad \begin{array}{ccc} Gr_e(R_n) & \xrightarrow{\varphi_n} & Gr_{e^*}(R_n) \\ N & \longrightarrow & N^0 \end{array}$$

where $(e_1, e_2)^* := (n - e_2, n - e_1)$ and $N^0 := \{f \in R_n^* \mid f(v) = 0 \ \forall v \in N\}$ by using the identification $R_n \simeq R_n^*$. The torus T acts on R_n^* with contragredient action and this gives an action on $Gr_{e^*}(R_n^*)$. Under the identification $R_n \simeq R_n^*$ these two actions differ by a character and so the

identification $Gr_{\mathbf{e}}(R_n^*) \simeq Gr_{\mathbf{e}}(R_n)$ is T -equivariant. It is an easy check that isomorphism (8) is T -equivariant and involutive.

2.3. The coefficient quiver of R_n . Following [23] we associate to a Q -representation M with standard basis $\mathcal{B}$ a quiver $\Gamma(M, \mathcal{B})$ called the coefficient quiver of M in the basis $\mathcal{B}$. By definition $\Gamma(M, \mathcal{B})$ has the elements of $\mathcal{B}$ as vertices and there is an arrow labeled by a (resp. b) between two vertices v and v' if the coefficient of v' in $m_a(v)$ (resp. $m_b(v)$) is non-zero. We call $\Gamma(M, \mathcal{B})$ the coefficient quiver of M in the basis $\mathcal{B}$. The following are the coefficient quivers of some indecomposable Q -representations for $n = 4$ in the standard basis. We denote by $k^{(i)}$ the vertex corresponding to $v_k^{(i)}$, $i = 1, 2$.

$$\begin{aligned} \Gamma(P_4) = & \begin{array}{ccccccc} & 1^{(1)} & & 2^{(1)} & & 3^{(1)} & & 4^{(1)} \\ & \swarrow a & \searrow b & \swarrow a & \searrow b & \swarrow a & \searrow b & \swarrow a & \searrow b \\ 1^{(2)} & & 2^{(2)} & & 3^{(2)} & & 4^{(2)} & & 5^{(2)} \end{array} \\ \Gamma(R_5(0)) = & \begin{array}{ccccccccc} & 1^{(1)} & & 2^{(1)} & & 3^{(1)} & & 4^{(1)} & & 5^{(1)} \\ & \swarrow a & \searrow b & \swarrow a & \searrow b & \swarrow a & \searrow b & \swarrow a & \searrow b & \swarrow a \\ 1^{(2)} & & 2^{(2)} & & 3^{(2)} & & 4^{(2)} & & 5^{(2)} & \end{array} \\ \Gamma(I_4) = & \begin{array}{ccccccc} & 1^{(1)} & & 2^{(1)} & & 3^{(1)} & & 4^{(1)} & & 5^{(1)} \\ & & \swarrow b & \searrow a & \swarrow b & \searrow a & \swarrow b & \searrow a & \swarrow b & \searrow a \\ & 1^{(2)} & & 2^{(2)} & & 3^{(2)} & & 4^{(2)} & & \end{array} \end{aligned}$$

In all these cases a one dimensional torus T acts on quiver Grassmannians and the fixed points of this action are in bijection with successor closed subquivers of the corresponding coefficient quiver.

Let us consider $\Gamma(R_n)$. In lemma 2.0.2 we have seen the torus T acts on X by $\lambda \cdot v_k^{(i)} = \lambda^{k-1} v_k^{(i)}$ ($i = 1, 2, k \in [1, n]$). For every $r \geq 1$ there exists a unique regular sub-representation of R_n isomorphic to R_r and it has the following property:

$$(9) \quad \text{if } N \in X \text{ is such that } \lim_{\lambda \rightarrow 0} t_\lambda N = R_r \text{ then } N = R_r.$$

Indeed this sub-representation is coordinate and lies in the extreme right hand side of $\Gamma(R_n)$. In particular the standard basis elements which generate R_r have maximal weights.

Roughly speaking the flow for $\lambda \rightarrow 0$ goes from right to left in $\Gamma(R_n)$. For example the line $\langle v_1^{(2)} + v_2^{(2)} \rangle$ generated by the vector $v_1^{(2)} + v_2^{(2)}$ goes to the line generated by $v_1^{(2)}$ as follows:

$$\lim_{\lambda \rightarrow 0} \lambda \cdot (\langle v_1^{(2)} + v_2^{(2)} \rangle) = \lim_{\lambda \rightarrow 0} (\langle v_1^{(2)} + \lambda v_2^{(2)} \rangle) = \langle v_1^{(2)} \rangle.$$

2.4. Action of the torus on Hom-spaces. Let L and L' be indecomposable Q -representations. The torus $T = \{t_\lambda : \lambda \in \mathbf{C}^*\}$ acts on the quiver Grassmannians associated with them. The action of T naturally extends to the vector space $Hom(L, L')$ as follows: for $f \in Hom(L, L')$, $(t_\lambda f)(l) := t_\lambda f(t_{\lambda^{-1}} l)$. Following [10] we endow the vector space $Hom(L, L')$ with a distinguished basis. Since $Hom(-, -)$ is additive, we assume that both L and L' are indecomposable. Let Γ_L and $\Gamma_{L'}$ denote the coefficient quiver of L and L' respectively. We consider the set (L, L') of triples $(\gamma, \alpha, \gamma')$




FIGURE 1. An element of the standard basis of $\text{Hom}(R_3, R_3)$: the circles (resp. bullets) highlight a predecessor (successor) closed subquiver γ (resp. γ') of $\Gamma(R_3)$. The dotted arrows show the corresponding $f_{\gamma\gamma'}$

such that γ is a connected *predecessor closed* subquiver of Γ_L (i.e. for every vertex $v \in \gamma$, every arrow $c : v' \rightarrow v$ with target v belongs to γ) and γ' is a successor closed subquiver of $\Gamma_{L'}$ and $\alpha : \gamma_0 \rightarrow \gamma'_0$ is a bijective map from the set γ_0 of vertices of γ to set γ'_0 of vertices of γ' such that for every arrow $v_k^{(1)} \xrightarrow{a} v_k^{(2)}$ (resp. $v_k^{(1)} \xrightarrow{b} v_{k+1}^{(2)}$) of γ there is an arrow $\alpha(v_k^{(1)}) \xrightarrow{a} \alpha(v_k^{(2)})$ (resp. $\alpha(v_k^{(1)}) \xrightarrow{b} \alpha(v_{k+1}^{(2)})$) of γ' with the same label. The next proposition is a special case of [10] and we hence avoid to repeat the proof (which is quite simple in this case).

Proposition 2.0.2. *We consider the map*

$$B : (L, L') \rightarrow \text{Hom}(L, L') : (\gamma, \alpha, \gamma') \mapsto f_{\gamma\gamma'}$$

which associates to $(\gamma, \alpha, \gamma')$ the homomorphism

$$f_{\gamma\gamma'}(v) = \begin{cases} \alpha(v) & \text{if } v \in \gamma_0, \\ 0 & \text{otherwise.} \end{cases}$$

The image of B is a basis of $\text{Hom}(L, L')$ which we call standard.

Figure 1 illustrates proposition 2.0.2. The torus T acts diagonally on the elements of the standard basis of $\text{Hom}(L, L')$ as follows: if $(\gamma, \alpha, \gamma') \in (L, L')$ and the vertices of γ have consecutive weights $k, k+1, \dots$ and the vertices of γ' have consecutive weights $k', k'+1, \dots$ then

$$t_\lambda f_{\gamma\gamma'} = \lambda^{k-k'} f_{\gamma\gamma'}$$

and we say that $f_{\gamma\gamma'}$ has weight $k - k'$. We say that an element $f_{\gamma\gamma'}$ has *positive weight* if $k - k' > 0$ and it has *negative weight* if $k - k' < 0$. We denote by $\text{Hom}(L, L')^+$ the vector subspace of $\text{Hom}(L, L')$ spanned by standard basis elements with positive weight. As an application of proposition 2.0.2 we compute the dimension of the Hom-spaces between indecomposable Q -representations. It is known (and not difficult to prove by using proposition 2.0.2) that $\text{Hom}(R_s, P_l) = \text{Hom}(I_s, P_l) = \text{Hom}(I_s, R_l) = 0$ for all $s, l \geq 0$. We hence consider the remaining cases.

Lemma 2.0.3. *For every $l, s \geq 0$ we have:*

$$(10) \quad \dim \operatorname{Hom}(P_s, P_l) = [l - s + 1]_+$$

$$(11) \quad \dim \operatorname{Hom}(P_s, R_l) = l$$

$$(12) \quad \dim \operatorname{Hom}(P_s, I_l) = l + s$$

$$(13) \quad \dim \operatorname{Hom}(R_s, R_l) = \min(s, l)$$

$$(14) \quad \dim \operatorname{Hom}(R_s, I_l) = s$$

$$(15) \quad \dim \operatorname{Hom}(I_s, I_l) = [s - l + 1]_+$$

where $[b]_+ := \max(b, 0)$.

We conclude this section by pointing out that the action of the torus T on $\operatorname{Hom}(L, L')$ affords an action of the torus on the space $\operatorname{Ext}^1(L, L')$. With respect to this action long exact sequences in cohomology are T -equivariant.

2.5. Stratification of X . Every sub-representation N of R_n is of the form $N = P \oplus R_r$ where P is preprojective and R_r is an indecomposable (possibly zero) regular Q -representation, for some $r \geq 0$. Similarly the quotient $R_n/N = R_{r'} \oplus I$ where $R_{r'}$ is regular indecomposable (possibly zero) and I is preinjective. We hence can give the following definition.

Definition 2.0.1. *Let $N \in X$ with $N = P \oplus R_r$ and $R_n/N = R_{r'} \oplus I$ with P preprojective, I preinjective and some $r, r' \geq 0$. We define the integer:*

$$K_N = K_N(X) := \min(r, r')$$

It is easy to see that $K_N = \dim \operatorname{Ext}^1(N, R_n/N)$. Indeed

$$\dim \operatorname{Ext}^1(N, R_n/N) = \dim \operatorname{Ext}^1(R_r, R_{r'}) = \dim \operatorname{Hom}(R_r, R_{r'}) = \min(r, r')$$

where in the first equality we use the well-known fact that $\operatorname{Ext}^1(R, I) = \operatorname{Ext}^1(P, R) = \operatorname{Ext}^1(P, I) = 0$ for every preprojective P , regular R and preinjective I Q -representations; in the second equality we use the AR-formula (see e.g. [1]); in the last equality we use (13). In particular it is known that

$$\langle \dim N, \dim N' \rangle = \dim \operatorname{Hom}(N, N') - \dim \operatorname{Ext}^1(N, N')$$

and in view of (6) we get

$$(16) \quad \dim T_N = \langle \mathbf{e}, n\delta - \mathbf{e} \rangle + K_N.$$

Equation (16) implies that a point $N \in X$ is smooth if and only if either N or M_n/N do not have a regular direct summand.

The next theorem provides a stratification of X and it is essential for our proof of the existence of a cellular decomposition of X . Let us define the strata.

Definition 2.0.2. *For every integer $k \geq 0$ define the set:*

$$X_k = X_k(X) := \{N \in X \mid K_N \geq k\}$$

where K_N is given in definition 2.0.1.

Theorem 2.1. (1) *The set X_{k+1} is a closed T -stable subvariety of X_k . Moreover there is a T -equivariant isomorphism:*

$$(17) \quad X_k \simeq \operatorname{Gr}_{(e_1-k, e_2-k)}(R_{n-2k});$$

- (2) The sets X_k provide a stratification of X
- (18)
$$X = X_0 \supset X_1 \supset \cdots \supset X_s$$
where $s = \min(e_1, n - e_2)$;
- (3) The variety $X_k \setminus X_{k+1}$ is smooth (inside X_k) and
- $$\forall N \in X_k \setminus X_{k+1}, \quad \lim_{\lambda \rightarrow 0} t_\lambda N \in X_k \setminus X_{k+1}$$

Proof. We consider the two subvarieties of X : for $k \in [0, e_1]$

$$(19) \quad X'_k(X) := \{N \in Gr_{\mathbf{e}}(R_n) \mid R_k \text{ is a subrepresentation of } N\}$$

and for $k \in [0, n - e_2]$

$$(20) \quad X''_k(X) := \{N \in Gr_{\mathbf{e}}(R_n) \mid R_k \text{ is a quotient of } R_n/N\}.$$

It follows from the definitions that:

$$(21) \quad X_k = X'_k \cap X''_k.$$

We now collect some properties of X'_k and X''_k .

Lemma 2.1.1. (1) The isomorphism φ_n defined in (8) induces an isomorphism

$$X'_k(Gr_{\mathbf{e}}(R_n)) \simeq X'_k(Gr_{\mathbf{e}^*}(R_n))$$

and hence also

$$X''_k(Gr_{\mathbf{e}}(R_n)) \simeq X''_k(Gr_{\mathbf{e}^*}(R_n)).$$

- (2) For every $k \in [0, e_1]$, $X'_k(X)$ is a T -stable subvariety of X . For every $N \in X'_k(X)$, $\lim_{\lambda \rightarrow 0} t_\lambda N \in X'_k(X)$. There is a T -equivariant isomorphism

$$(22) \quad X'_k(X) \simeq Gr_{\mathbf{e}-k\delta}(R_{n-k})$$

- (3) For every $k \in [0, n - e_2]$, $X''_k(X)$ is a T -stable subvariety of X . For every $N \in X''_k(X)$, $\lim_{\lambda \rightarrow 0} t_\lambda N \in X''_k(X)$. There is a T -equivariant isomorphism

$$(23) \quad X''_k(X) \simeq Gr_{\mathbf{e}}(R_{n-k})$$

Proof. Part (1) follows by straightforward check. Part (3) follows from part (1) by using the isomorphism (8). Part (2) follows from property (9) of R_k . It remains to check (22). We consider the map:

$$\begin{aligned} \{N \in Gr_{\mathbf{e}}(R_n) \mid R_k \leq N\} &\longrightarrow Gr_{\mathbf{e}-k\delta}(R_{n-k}) \\ N &\longmapsto N/R_k \end{aligned}$$

Since R_n is uniserial, there is a canonical short exact sequence:

$$(24) \quad 0 \longrightarrow R_k \xrightarrow{\iota_k} R_n \xrightarrow{\pi_{n-k}} R_{n-k} \longrightarrow 0$$

and hence the map is well defined and bijective. $\square$

We now conclude the proof of the theorem. Everything follows from lemma 2.1.1 by (21) except the smoothness of $X_k \setminus X_{k+1}$. To prove smoothness notice that it is sufficient to prove smoothness of $X_0 \setminus X_1$ by (17), since $K_N(X_k) = K_N(X) - k$. But by (16) $X_0 \setminus X_1$ is the smooth locus of X . $\square$

Corollary 2.1.1. *The variety $X = \text{Gr}_{\mathbf{e}}(R_n)$ is smooth if and only if $e_1 = 0$ (in which case $X \simeq \text{Gr}_{e_2}(k^n)$) or $e_2 = n$ (in which case $X \simeq \text{Gr}_{e_1}(k^n)$).*

Proof. By theorem 2.1, X is smooth if and only if $X = X_0$ if and only if $s = \min(e_1, n - e_2) = 0$. $\square$

Corollary 2.1.2. *The quiver Grassmannian $\text{Gr}_{(e_1, e_2)}$ has dimension:*

$$\dim \text{Gr}_{(e_1, e_2)} = \langle \mathbf{e}, n\delta - \mathbf{e} \rangle = (e_2 - e_1)(n - (e_2 - e_1))$$

Proof. It follows from (16). $\square$

2.6. Cellular decomposition of X . In this section we provide a cellular decomposition of $X = \text{Gr}_{\mathbf{e}}(R_n)$. Following [3] (see also [9]), for every fixed point $L \in X^T$ we consider its *attracting set* defined as follows:

$$(25) \quad X_L := \{N \in X \mid \lim_{\lambda \rightarrow 0} t_\lambda N = L\}.$$


In particular $L \in X_L$ for every $L \in X^T$.

Theorem 2.2. *For every $L \in X^T$ the corresponding attracting set X_L is an affine space and $X_L \simeq T_L(X_L) \simeq T_L^+(X) := \text{Hom}(L, R_n/L)^+$. Moreover*

$$X = \cup_{L \in X^T} X_L$$

Proof. Bialynicki–Birula’s results on cellular decomposition of a projective variety X [3] continues to hold if the variety is smooth but only quasi-projective provided that the action of the torus is such that $\lim_{\lambda \rightarrow 0} t_\lambda N$ belongs to X for every $N \in X$. This is an easy consequence of Hironaka’s resolution of singularities. Since of theorem 2.1 we apply this to $X_0 \setminus X_1$ and we get the result. $\square$

2.7. Description of the cells. In this section we describe the cell X_L associated with every $L \in X^T$ (see theorem 2.2). If $L \simeq P_r$ is an indecomposable preprojective subrepresentation of R_n of dimension $(r, r+1)$, we write $L := {}_k(P_{e_1})$ if L is generated by $v_k^{(1)}, v_{k+1}^{(1)}, \dots, v_{k+r-1}^{(1)}$ and $v_k^{(2)}, v_{k+1}^{(2)}, \dots, v_{k+r}^{(2)}$. For example the following figure shows the subrepresentation ${}_2(P_1)$ of R_5 :



Theorem 2.3. (1) *If $L \in X^T$ is indecomposable then $X_L = \mathcal{U}L$. In particular if $L = {}_k(P_{e_1})$ then $\dim X_L = n - k$;*

(2) *If $L = L' \oplus L''$ with $\text{Hom}(L', L'')^+ = \text{Hom}(L', L'')$ then*

$$(26) \quad \dim X_L = \dim X_{L'} + \dim X_{L''} - \langle \dim L', \dim L'' \rangle$$

Proof. We prove part (1). The unipotent group $\mathcal{U}$ defined in lemma 2.0.2 is a subgroup of dimension $n - 1$ of the group of $n \times n$ unipotent triangular matrices. Let L be an indecomposable sub-representation of R_n of dimension vector $\mathbf{e}$. If $L = R_r$ then $\mathbf{e} = (r, r)$ and $L_1 = L_2$ is the vector subspace of k^n spanned by the last r standard basis vectors. In particular $AL = L$ for every $A \in \mathcal{U}$. On the other hand we have already noticed that regular sub-representations have property (9) and hence $X_L = \{L\}$ and we get $X_L = \mathcal{U}L$ if L is regular. Let L be an indecomposable preprojective sub-representation

of R_n . Then $e_2 = e_1 + 1$ and we assume that L_1 is spanned by consecutive standard basis vectors $\{v_k, v_{k+1}, \dots, v_{k+e_1-1}\}$ and L_2 is spanned by L_1 and v_{k+e_1} . The following easy lemma says that $\dim \mathcal{U}L = n - k$.

Lemma 2.3.1. *Let W be a vector subspace of k^n spanned by some basis vectors $\{v_i : i \geq k\}$ and v_k is in this set. Then the stabilizer $\text{Stab}_{\mathcal{U}}(W) = \text{Stab}_{\mathcal{U}}(< v_k >)$. In particular the orbit $\mathcal{U}W$ has dimension $n - k$.*

Proof. We prove that $\text{Stab}_{\mathcal{U}}(W) \supset \text{Stab}_{\mathcal{U}}(< v_k >)$ being the other inclusion clear. The columns $\mathbf{a}_1, \dots, \mathbf{a}_n$ of a matrix $A \in \mathcal{U}$ satisfy the relation: $\mathbf{a}_l = J_n(0)\mathbf{a}_{l-1}$ for every $l = 2, 3, \dots, n$. Let $A \in \text{Stab}_{\mathcal{U}}(< v_k >)$ and $Av_k = \beta v_k$ for some non-zero scalar β . Let $v := \sum_{i \geq 0} b_i v_{k+i}$ be an element of W ; we get:

$$Av = \sum_{i \geq 0} b_i Av_{k+i} = \sum_{i \geq 0} b_i \mathbf{a}_{k+i} = \sum_{i \geq 0} b_i J^i \mathbf{a}_k = \sum_{i \geq 0} b_i J^i Av_k = \sum_i b_i \beta J^i v_k$$

which belongs to W and hence $A \in \text{Stab}_{\mathcal{U}}(W)$. In particular $\mathcal{U}W = \mathcal{U}(< v_k >)$ has dimension $n - k$. $\square$

We now prove that $\dim X_L = n - k$ and since $\mathcal{U}L \subset X_L$ we get the equality. By theorem 2.2 we have to compute $\dim T_L^+(X) = \dim \text{Hom}(L, R_n/L)^+$. Since L is indecomposable the quotient R_n/L is the direct sum of at most two indecomposables as follows

$$R_n/L = R_{k-1} \oplus I_t$$

where $t = n - 1 - k - e_1$. We have:

$$\text{Hom}(L, R_n/L)^+ = \text{Hom}(P_{e_1}, I_t).$$

In view of (12), $\dim(\text{Hom}(L, R_n/L)^+) = e_1 + t = n - k$ and we are done.

We now prove part (2). There are short exact sequences:

$$0 \longrightarrow L' \longrightarrow R_n/L'' \longrightarrow R_n/(L' \oplus L'') \longrightarrow 0$$

and

$$0 \longrightarrow L'' \longrightarrow R_n/L' \longrightarrow R_n/(L' \oplus L'') \longrightarrow 0$$

The torus acts on Hom-spaces and on Ext-spaces between T -fixed points as explained in section 2.4. Moreover these spaces split into the subspaces spanned by standard basis vectors with positive and negative weights respectively. We apply the functors $\text{Hom}(L'', -)$ and $\text{Hom}(L', -)$ to the previous short exact sequences, we take the positive part and we get the following exact sequences:

$$0 \longrightarrow 0 \longrightarrow T_{L''}(X_{L''}) \longrightarrow \text{Hom}(L'', R_n/L) \longrightarrow 0$$

and

$$0 \rightrightarrows \text{Hom}(L', L'') \rightrightarrows T_{L'}(X_{L'}) \rightrightarrows \text{Hom}(L', R_n/L) \rightrightarrows \text{Ext}^1(L', L'')^+ \rightrightarrows 0$$

where we use the following equalities:

$$(27) \quad \text{Ext}^1(L'', L')^+ = 0 = \text{Ext}^1(L', R_n/L)^+$$

$$(28) \quad \text{Ext}^1(L', L'')^+ = \text{Ext}^1(L', L'')$$

This follows by the fact that if $\text{Hom}(M, L)^+ = 0$ then $\text{Ext}^1(M, L)^+ = 0$; indeed one can always take a minimal T -equivariant injective resolution of L and apply the functor $\text{Hom}(M, -)$. By summing up we get the short exact sequence:

$$0 \rightarrow \text{Hom}(L', L'') \rightarrow T_{L'}(X_{L'}) \oplus T_{L''}(X_{L''}) \rightarrow T_L(X_L) \rightarrow \text{Ext}^1(L', L'') \rightarrow 0$$

Since for every fixed point L , $\dim X_L = \dim T_L(X_L)$ we get (26). $\square$

2.8. Betti numbers. We now use the results of the previous sections in order to compute the Betti numbers of $X = \text{Gr}_{\mathbf{e}}(R_n)$. Since X has a cellular decomposition (theorem 2.2) the odd cohomology spaces are zero and the $2i$ -th Betti number $b_{2i} = b_{2i}(X) := \dim H^{2i}(X)$ equals the number of cells of dimension i .

Before stating the main result of this section we start with the special case $e_2 = e_1 + 1$.

Theorem 2.4. *Let $X = \text{Gr}_{(e_1, e_1+1)}(R_n)$. The even Betti numbers of X are the following:*

$$(29) \quad b_{2i} = \begin{cases} i+1 & \text{if } 0 \leq i \leq s, \\ s+1 & \text{if } s \leq i \leq n-1-s, \\ n-i & \text{if } n-1-s \leq i \leq n-1 \end{cases}$$

where $s = \min(e_1, n-1-e_1)$.

The Poincaré polynomial $P_X(t) := \sum_{i=0}^{\dim X} b_i t^i$ of X equals:

$$(30) \quad P_X(t^{1/2}) = \left(\frac{t^{e_1+1} - 1}{t - 1} \right) \left(\frac{t^{n-e_1} - 1}{t - 1} \right)$$

Proof. For $k \in [0, e_1]$ let us consider the variety X'_k defined in (19) and the difference

$$(31) \quad Y'_k = Y'_k(X) := X'_k(X) \setminus X'_{k+1}(X).$$

By definition $N \in Y_k$ if $N = R_k \oplus P$ for some preprojective P . By lemma 2.1.1 we have

$$(32) \quad \dim H^i(X) = \sum_{k=0}^{e_1} \dim H_c^i(Y'_k)$$

where $H_c^i(Y'_k)$ is the i -th cohomology space of Y'_k with compact support. Moreover $\dim H_c^i(Y'_k) = 0$ for i odd and

$$(33) \quad \dim H_c^{2i}(Y'_k) = \begin{cases} 1 & \text{if } i \in [e_1 - k, n - k - 1] \\ 0 & \text{otherwise.} \end{cases}$$

and hence (29) follows from (32). It remains to prove (33). It is easy to see that

$$(34) \quad Y'_k(X) \simeq Y'_0(\text{Gr}_{e_1-k, e_2-k}(R_{n-k})).$$

The elements of $Y'_0(\text{Gr}_{(e_1, e_1+1)}(R_n))$ are all the indecomposable preprojective subrepresentations of R_n of dimension $(e_1, e_1 + 1)$ and hence they are all isomorphic to P_{e_1} . We denote by ${}_i(P_{e_1})$ the fixed point of $Y'_0(\text{Gr}_{(e_1, e_1+1)}(R_n))$ generated by the vectors $\{v_i^{(1)}, v_{i+1}^{(1)}, \dots, v_{i+e_1-1}^{(1)}\}$ and $\{v_i^{(2)}, v_{i+1}^{(2)}, \dots, v_{i+e_1}^{(2)}\}$ for $i \in [1, n - e_1]$. By theorem 2.3 $\dim X_{i(P_{e_1})} = n - i \in [e_1, n - 1]$ and

hence for every $i \in [e_1, n-1]$ there is a unique cell of dimension i and we get (33). $\square$

Notice that (30) can be written as follows:

$$P_X(t) = P_{Gr_1(e_2)}(t)P_{Gr_1(n-e_1)}(t)$$

where $Gr_s(t)$ denotes the Grassmannians of s -dimensional vector subspaces of a t -dimensional vector space. Surprisingly this turns out to be a general fact and it is the main result of this section.

Theorem 2.5. *Let $X = Gr_{(e_1, e_2)}(R_n)$. The Poincaré polynomial $P_X(t)$ of X equals*

$$(35) \quad P_X(t) = P_{Gr_{(e_2-e_1)}(e_2)}(t)P_{Gr_{(e_2-e_1)}(n-e_1)}(t).$$

Proof. We proceed by induction on $n \geq e_1 \geq 0$. For $e_1 = 0$, $X \simeq Gr_{e_2}(n)$ and (35) follows. Let $1 \leq e_1 \leq e_2$. The variety X can be decomposed in

$$X = Y'_0(X) \cup X'_1(X)$$

where $Y'_0 = Y'_0(X)$ consists of all preprojective subrepresentations of R_n in X and X'_1 consists of subrepresentations of R_n in X having a non-zero regular subrepresentation (see (19) and (31)). Moreover, by (22), $X'_1(X) \simeq Gr_{(e_1-1, e_2-1)}(R_{n-1})$ and we hence have:

$$(36) \quad P_X(t) = P_{Y'_0}(t) + P_{Gr_{(e_1-1, e_2-1)}(R_{n-1})}(t).$$

By inductive hypothesis the equality (35) holds if and only if the following equality holds:

$$(37) \quad P_{Y'_0}(t) = \left(P_{Gr_{(e_2-e_1)}(e_2)}(t) - P_{Gr_{(e_2-e_1)}(e_2-1)}(t) \right) P_{Gr_{(e_2-e_1)}(n-e_1)}(t).$$

We hence prove (37). We make the following choice: we fix a basis $v_1 \cdots v_s$ of an s -dimensional vector space and we let the torus act on $Gr_t(s)$ by $t_\lambda v_i = \lambda^i v_i$. We consider the vector subspace of k^{e_2} generated by $v_2, \dots, v_{e_2}$ and the corresponding embedding $Gr_{(e_2-e_1)}(e_2-1) \subset Gr_{(e_2-e_1)}(e_2)$. With this choices the difference $Gr_{(e_2-e_1)}(e_2) \setminus Gr_{(e_2-e_1)}(e_2-1)$ is T -stable and for every point W of it, $\lim_{\lambda \rightarrow 0} t_\lambda W$ still belongs to it. The right hand side of (37) is the Poincaré polynomial (with respect to the cohomology with compact support) of the smooth projective variety:

$$G := (Gr_{(e_2-e_1)}(e_2) \setminus Gr_{(e_2-e_1)}(e_2-1)) \times Gr_{(e_2-e_1)}(n-e_1)$$

The one-dimensional torus T acts on G and the attracting sets of the T -fixed points form a cellular decomposition of G . We prove that there exists a bijection between the cells of Y'_0 of dimension k and the cells of G of dimension k .

A point of Y'_0 is a direct sum of precisely $(e_2 - e_1)$ preprojective subrepresentations of R_n (this follows by considering their dimension vector). The T -fixed points have the form:

$$L := k_1(P_{r_1}) \oplus k_2(P_{r_2}) \oplus \cdots \oplus k_{(e_2-e_1)}(P_{r_{(e_2-e_1)}})$$

where $r_1 + \cdots + r_{e_2-e_1} = e_1$, $r_i \geq 0$ and $k(P_r)$ denotes the unique indecomposable preprojective subrepresentation of R_n of dimension vector $(r, r+1)$

generated by $v_k^{(1)}, v_{k+1}^{(1)}, \dots, v_{k+r-1}^{(1)}$ and $v_k^{(2)}, v_{k+1}^{(2)}, \dots, v_{k+r}^{(2)}$. In view of theorem 2.3 it is easy to see that the dimension of the attracting cell of L equals

$$\dim X_L = n(e_2 - e_1) - \sum_{i=1}^{e_2-e_1} k_i - (e_2 - e_1)^2 + \sum_{i=1}^{e_2-e_1} \sum_{j=1}^i (r_j - r_i + 1).$$

We consider the set $\alpha(\mathbf{e}, k, n)$ which parametrizes the T -fixed points of Y'_0 whose attracting set has dimension k , i.e.

$$\begin{aligned} \alpha(\mathbf{e}, k, n) := & \{(k_1, k_2, \dots, k_{e_2-e_1}, r_1, r_2, \dots, r_{e_2-e_1}) \mid \\ & 1 \leq k_1 \leq k_1 + r_1 < k_2 \leq k_2 + r_2 < \dots < k_{e_2-e_1} \leq k_{e_2-e_1} + r_{e_2-e_1} \leq n; \\ & r_1 + r_2 + \dots + r_{e_2-e_1} = e_1, r_i \geq 0; \\ & n(e_2 - e_1) - \sum_{i=1}^{e_2-e_1} k_i - (e_2 - e_1)^2 + \sum_{i=1}^{e_2-e_1} \sum_{j=1}^i (r_j - r_i + 1) = k\}. \end{aligned}$$

On the other hand we consider the T -fixed points of G and their attracting sets. The T -fixed points of $Gr_t(s)$ consist of coordinate vector subspaces of dimension t and they are naturally parametrized by tuples $(a_1, \dots, a_t)$ of integers $1 \leq a_1 < \dots < a_t \leq s$. The corresponding cell $\mathcal{O}_{(a_1, \dots, a_t)}$ has dimension

$$s - a_1 - (t - 1) + s - a_2 - (t - 2) + \dots + s - a_t = ts - \sum_{i=1}^t a_i - \sum_{i=1}^{t-1} i.$$

The T -fixed points of $Gr_{(e_2-e_1)}(e_2) \setminus Gr_{(e_2-e_1)}(e_2 - 1)$ are the coordinate vector subspaces of k^{e_2} containing v_1 . The following set hence parametrizes the cells of G of dimension k :

$$\begin{aligned} \beta(\mathbf{e}, k, n) := & \{(a_1, a_2, \dots, a_{e_2-e_1}, b_2, b_3, \dots, b_{e_2-e_1}) \mid \\ & 1 \leq a_1 < a_2 < \dots < a_{e_2-e_1} \leq n - e_1; \\ & 2 \leq b_2 < \dots < b_{e_2-e_1} \leq e_2; \\ & n(e_2 - e_1) - \sum_{i=1}^{e_2-e_1} a_i - \sum_{i=2}^{e_2-e_1} b_i + (e_2 - e_1 - 1) = k\}. \end{aligned}$$

We consider the map

$$(38) \quad a_1 = k_1$$

$$(39) \quad a_i = k_i - r_1 - r_2 - \dots - r_{i-1}$$

$$(40) \quad b_i = \sum_{j=0}^{i-2} r_{(e_2-e_1-j)} + i$$

for $i \in [2, e_2 - e_1]$. It is straightforward to verify that this map is a bijection between $\alpha(\mathbf{e}, k, n)$ and $\beta(\mathbf{e}, k, n)$. It follows that Y'_0 and G have the same Betti numbers and hence (37) follows. $\square$

Corollary 2.5.1. *The Poincaré polynomial of a quiver Grassmannian associated with the indecomposable preprojective P_n and the indecomposable preinjective I_n ($n \geq 0$) are respectively the following:*

$$(41) \quad P_{Gr_{\mathbf{e}}(P_n)}(t) = P_{Gr_{e_1}(e_2-1)}(t) P_{Gr_{(e_2-e_1)}(n+1-e_1)}(t)$$

$$(42) \quad P_{Gr_{\mathbf{e}}(I_n)}(t) = P_{Gr_{e_1}(e_2+1)}(t) P_{Gr_{(e_2-e_1)}(n-e_1)}(t)$$

Proof. The equality (42) follows from (41) by the isomorphism:

$$Gr_{(e_1, e_2)}(I_n) \simeq Gr_{(n-e_2, n+1-e_1)}(P_n).$$

We hence prove (41). As in the proof of theorem 2.5, let $Y'_0 = Y'_0(Gr_{\mathbf{e}}(R_{n+1}))$ be the subvariety of $Gr_{\mathbf{e}}(R_{n+1})$ of all preprojective subrepresentations and let

$$G' := Gr_{(e_2-e_1)}(e_2) \setminus Gr_{(e_2-e_1)}(e_2-1)$$

with the convention that $Gr_{(e_2-e_1)}(e_2-1)$ consists of all the elements of $Gr_{(e_2-e_1)}(e_2)$ not containing the first basis vector (as in the proof of theorem 2.5). In view of (37) it is sufficient to prove the following equalities:

$$(43) \quad P_{G'}(t) = t^{2e_1} P_{Gr_{(e_2-e_1-1)}(e_2-1)}(t),$$

$$(44) \quad P_{Y'_0}(t) = t^{2e_1} P_{Gr_{\mathbf{e}}(P_n)}(t).$$

The proof of (43) is similar to the proof of theorem 2.5: there is an obvious bijection between the cells of G' of dimension k and the cells of $Gr_{(e_2-e_1)}(e_2-1)$ of dimension $k-e_1$. Let us prove (44). Let $L \in Y'_0$. Then L is a sum of preprojective subrepresentations of R_{n+1} and L is a subrepresentation of $P_n \leq R_{n+1}$. After a look at the quotients R_{n+1}/L and P_n/L and using lemma 2.0.3 one gets the following:

$$\dim Hom(L, R_{n+1}/L)^+ = \dim Hom(L, P_n/L)^+ + e_1$$

and hence the cells of Y'_0 of dimension k are in bijection with the cells of $Gr_{\mathbf{e}}(P_n)$ of dimension $k-e_1$ and (44) holds. $\square$

Corollary 2.5.2.

$$(45) \quad \chi(Gr_{\mathbf{e}}(R_n)) = \binom{e_2}{e_1} \binom{n-e_1}{e_2-e_1}.$$

$$(46) \quad \chi(Gr_{\mathbf{e}}(P_n)) = \binom{e_2-1}{e_1} \binom{n+1-e_1}{e_2-e_1}$$

$$(47) \quad \chi(Gr_{\mathbf{e}}(I_n)) = \binom{e_2+1}{e_1} \binom{n-e_1}{e_2-e_1}$$

Proof. For a projective variety X , $\chi(X) = P_X(1)$ and $\chi(Gr_t(s)) = \binom{s}{t}$. $\square$

3. APPLICATION TO CLUSTER ALGEBRAS

To a finite quiver Q without loops and 2-cycles is associated a (coefficient-free) cluster algebra $\mathcal{A}_Q$ ([13], see also [15] and [20] for excellent surveys). This is a $\mathbb{Z}$ -subalgebra of the field of rational functions in n ($=$ number of vertices of Q) variables generated by its *cluster variables*. The *cluster monomials* are monomials in cluster variables belonging to the same cluster. A canonical basis of $\mathcal{A}_Q$ is a $\mathbb{Z}$ -basis $\mathbf{B}$ of it such that the positive linear combinations of elements of $\mathbf{B}$ coincide with the semiring of positive elements of $\mathcal{A}_Q$, i.e. the elements which are positive Laurent polynomials in every cluster of $\mathcal{A}_Q$ (see [24]). The existence of such a basis has been proved only in a few cases:

- when Q is the Kronecker quiver, the canonical basis has been found by Sherman and Zelevinsky [24] (for every choice of the coefficients) and consists of cluster monomials together with extra elements $\{z_n : n \geq 1\}$;
- when Q is of type $A_2^{(1)}$, the canonical basis of $\mathcal{A}_Q$ has been found in [18] (for every choice of the coefficients) and consists of cluster monomials together with some extra elements $\{u_n : n \geq 1\}$ possibly multiplied with some cluster variables.

Note that under the Caldero–Keller bijection, cluster monomials correspond to rigid Q –representations and the quiver Grassmannians associated with rigid representations are smooth. In the next two sections we give a geometric realization of the extra elements $\{z_n\}$, $\{u_n\}$.

3.1. Type $A_1^{(1)}$. Let $\mathcal{F} = \mathbf{Q}(x_1, x_2)$ be the field of rational functions in two independent variables x_1 and x_2 with rational coefficients. We define recursively elements $\{x_k \mid k \in \mathbb{Z}\}$ of $\mathcal{F}$ by:

$$x_k x_{k+2} = x_{k+1}^2 + 1, \quad k \in \mathbb{Z}.$$

Let $\mathcal{A}$ be the $\mathbb{Z}$ –subalgebra of $\mathcal{F}$ generated by the x_k ($k \in \mathbb{Z}$). By [24] the algebra $\mathcal{A}$ is the coefficient–free cluster algebra associated with the Kronecker quiver. The couples $\{x_k, x_{k+1}\}$, $k \in \mathbb{Z}$, are free generating sets of $\mathcal{F}$ and they are called the *clusters* of $\mathcal{A}$. Monomials $x_k^a x_{k+1}^b$, $a, b \geq 0$, $k \in \mathbb{Z}$, are called the *cluster monomials* of $\mathcal{A}$.

In [8], Caldero and Zelevinsky have defined the rational function:

$$s_n := CC(R_n)$$

for every $n \geq 0$, where R_n is a regular indecomposable Q –representation of dimension (n, n) . They have proved that the set

$$\mathcal{S} := \{\text{cluster monomials}\} \cup \{s_n : n \geq 1\}$$

is a $\mathbb{Z}$ –basis of $\mathcal{A}$.

The canonical basis $\mathbf{B}$ defined in the introduction and this basis are related by the following formula [8]:

$$(48) \quad z_n = s_n - s_{n-2}$$

for $n \geq 1$ and the convention that $s_{-k} = 0$ for $k > 0$.

Theorem 3.1. *The element z_n has the following Laurent expansion:*

$$(49) \quad z_n = \frac{\sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(R_n)^{Sm}) x_1^{2(n-e_2)} x_2^{2e_1}}{x_1^n x_2^n}$$

where $\text{Gr}_{\mathbf{e}}(R_n)^{Sm}$ denotes the smooth part of $\text{Gr}_{\mathbf{e}}(R_n)$.

Proof. In view of theorem 2.1, $\text{Gr}_{\mathbf{e}}(R_n)^{Sm} = X_0 \setminus X_1$ where $X_0 = \text{Gr}_{\mathbf{e}}(R_n)$ and $X_1 = \text{Gr}_{(e_1-1, e_2-1)}(R_{n-2})$ and hence:

$$\chi(\text{Gr}_{\mathbf{e}}(R_n)^{Sm}) = \chi(\text{Gr}_{\mathbf{e}}(R_n)) - \chi(\text{Gr}_{(e_1-1, e_2-1)}(R_{n-2})).$$

It is now easy to check that the right hand side of (49) satisfies (48). $\square$

3.2. Type $A_2^{(1)}$. We now briefly recall the construction of the canonical basis of cluster algebras of type $A_2^{(1)}$ from [18]. Let $\mathcal{F} = \mathbb{Q}(x_1, x_2, x_3)$ be the field of rational functions in three (commuting) independent variables x_1 , x_2 and x_3 with rational coefficients. Recursively define elements $x_m \in \mathcal{F}$ for $m \in \mathbb{Z}$ by the relations

$$(50) \quad x_m x_{m+3} = x_{m+1} x_{m+2} + 1.$$

Define also the elements $w, z \in \mathcal{F}$ by

$$(51) \quad w = \frac{x_1 + x_3}{x_2}$$

$$(52) \quad z = \frac{x_1 x_2 + x_2 x_3 + 1}{x_1 x_3}$$

The (coefficient-free) cluster algebra $\mathcal{A}$ of type $A_2^{(1)}$ is the $\mathbb{Z}$ -subalgebra of $\mathcal{F}$ generated by all the x_m , w and z (see also [13, Example 7.8]). This is the cluster algebra associated with the affine quiver Q_2 of type $A_2^{(1)}$ shown in figure 2. The elements x_m , $m \in \mathbb{Z}$, w and z are the cluster variables




FIGURE 2. The quiver of type $A_2^{(1)}$

of $\mathcal{A}$. The sets $\{x_m, x_{m+1}, x_{m+2}\}$, $\{x_{2m}, z, x_{2m+2}\}$ and $\{x_{2m-1}, w, x_{2m+1}\}$, $m \in \mathbb{Z}$, are the clusters of $\mathcal{A}$. The cluster monomials are monomials in cluster variables belonging to the same cluster. The exchange graph of $\mathcal{A}$ is the brick wall shown in figure 3: it has clusters as vertices and an edge between two vertices if the corresponding clusters share precisely two cluster variables. In this figure the cluster variables of a cluster label the regions surrounding its corresponding vertex. Define elements u_n , $n \geq 0$, of $\mathcal{F}$ by




FIGURE 3. The exchange graph of a cluster algebra of type $A_2^{(1)}$

the recursion:

$$(53) \quad \begin{aligned} u_0 &= 2 \\ u_1 &= zw - 2 \\ u_{n+1} &= u_1 u_n - u_{n-1} \quad n \geq 1 \end{aligned}$$

In [18] it is shown that the set

$$(54) \quad \mathbf{B} = \{\text{cluster monomials}\} \cup \{u_n w^k, u_n z^k : n \geq 1, k \geq 0\}$$

is a $\mathbb{Z}$ -basis of $\mathcal{A}$ such that the positive elements of $\mathcal{A}$ are precisely the non-negative linear combinations of elements of $\mathbf{B}$. We now realize the elements u_n as image of the Caldero–Chapoton map. Recall that for a representation M of the quiver Q_2 shown in figure 2 the Caldero–Chapoton map $CC(M)$ is the following (see [8]):

$$CC(M) = \frac{\sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(M)) x_1^{d_2+d_3-e_2-e_3} x_2^{d_3-e_3+e_1} x_3^{e_1+e_2}}{x_1^{d_1} x_2^{d_2} x_3^{d_3}}$$

where (d_1, d_2, d_3) is the dimension vector of M . For every $n \geq 1$ let $R_{n,2}$ be the indecomposable regular Q_2 -representation in an homogeneous tube, i.e.

$$R_{n,2} = \begin{array}{ccc} & k^n & \\ \nearrow & & \searrow \\ k^n & \xrightarrow{J_n(0)} & k^n \end{array}$$

where $J_n(0)$ denotes the $n \times n$ indecomposable nilpotent Jordan block.

Theorem 3.2. *For every $n \geq 1$:*

$$(55) \quad u_n = \frac{\sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(R_{n,2})^{Sm}) x_1^{2n-e_2-e_3} x_2^{n-e_3+e_1} x_3^{e_1+e_2}}{x_1^n x_2^n x_3^n}$$

where $\text{Gr}_{\mathbf{e}}(R_{n,2})^{Sm}$ denotes the smooth part of the quiver Grassmannian $\text{Gr}_{\mathbf{e}}(R_{n,2})$.

Proof. Let u'_n denote the right hand side of (55). We consider the fibration of projective varieties

$$\begin{aligned} f : \text{Gr}_{(e_1, e_2, e_3)}(R_{n,2}) &\longrightarrow \text{Gr}_{(e_1, e_3)}(R_n) \\ (N_1, N_2, N_3) &\longmapsto (N_1, N_3) \end{aligned}$$

whose fiber is $\text{Gr}_{(e_2-e_1)}(e_3-e_1)$. In particular

$$\text{Gr}_{(e_1, e_2, e_3)}(R_{n,2})^{Sm} = f^{-1}(\text{Gr}_{(e_1, e_3)}(R_n)^{Sm}).$$

Using this it is straightforward to check that

$$(56) \quad u'_n = z_n(x_1 w^{-1/2}, x_3 w^{-1/2})$$

where the right hand side means to compute (49) with $x_1 \mapsto x_1 w^{-1/2}$ and $x_2 \mapsto x_3 w^{-1/2}$ and $w^{-1/2}$ is a formal variable whose square is w^{-1} . By definition (4) we have

$$u'_1 = z'_1(x_1 w^{-1/2}, x_3 w^{-1/2}) = \frac{x_1^2 + x_3^2 + w}{x_1 x_2} = zw - 2$$

and $u'_{n+1} = u'_1 u'_n - u'_{n-1}$ and hence $u'_n = u_n$ for every $n \geq 1$. $\square$

4. CONCLUSIONS

The study of quiver Grassmannians from a geometric point of view has given interesting information about cluster algebras. After this study it is natural to define the following slight modification of the Caldero–Chapoton map: for $k \geq 0$

$$(57) \quad CC^{(k)}(M) := \frac{\sum_{\mathbf{e}} \chi(Gr_{\mathbf{e}}(M)^{(k)}) x_1^{2(d_2-e_2)} x_2^{2e_1}}{x_1^{d_1} x_2^{d_2}}$$

where

$$Gr_{\mathbf{e}}(M)^{(k)} := \{N \in Gr_{\mathbf{e}}(M) \mid \dim Ext^1(N, M/N) = k\}.$$

So $CC^{(0)}(M)$ is defined by the smooth part of the quiver Grassmannians associated with M . In particular for a rigid representation M we have $CC^{(0)}(M) = CC(M)$. Moreover it is easy to check that if $Ext^1(M, N) = Ext^1(N, M) = 0$ then $CC^0(M \oplus N) = CC^0(M)CC^0(N)$. In particular the elements $u_n w$ and $u_n z$ of the basis (54) are $CC^0(R_{n,2} \oplus W)$ and $CC^0(R_{n,2} \oplus Z)$ where W and Z are the indecomposable regular rigid Q_2 -representations.

It would be interesting to study the relation between the nice construction due to Dupont [12] of transverse quiver Grassmannians with the maps $CC^{(k)}$.

We hope that the technique developed here can help in the determination of a “canonical basis” of cluster algebras of higher rank which is still an open problem.

Acknowledgements. The first author thanks the organizers of the “Advanced School and Conference on Homological and Geometrical Methods in Representation Theory” which took place in Trieste (Italy) at the beginning of 2010 for the opportunity to present this project in front of several experts and to spend a fruitful time there.

We are grateful to Andrei Zelevinsky for having pointed out the formulas found by Szanto in [25].

We thank Grégoire Dupont for useful comments about transverse quiver Grassmannians.

REFERENCES

- [1] Ibrahim Assem, Daniel Simson, and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.
- [2] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky. Cluster algebras. III. Upper bounds and double Bruhat cells. *Duke Math. J.*, 126(1):1–52, 2005.
- [3] A. Białynicki-Birula. Some theorems on actions of algebraic groups. *Ann. of Math. (2)*, 98:480–497, 1973.
- [4] Philippe Caldero and Frédéric Chapoton. Cluster algebras as Hall algebras of quiver representations. *Comment. Math. Helv.*, 81(3):595–616, 2006.
- [5] Philippe Caldero and Bernhard Keller. From triangulated categories to cluster algebras. II. *Ann. Sci. École Norm. Sup. (4)*, 39(6):983–1009, 2006.
- [6] Philippe Caldero and Bernhard Keller. From triangulated categories to cluster algebras. *Invent. Math.*, 172(1):169–211, 2008.
- [7] Philippe Caldero and Markus Reineke. On the quiver Grassmannian in the acyclic case. *J. Pure Appl. Algebra*, 212(11):2369–2380, 2008.

- [8] Philippe Caldero and Andrei Zelevinsky. Laurent expansions in cluster algebras via quiver representations. *Mosc. Math. J.*, 6(3):411–429, 2006.
- [9] Neil Chriss and Victor Ginzburg. *Representation theory and complex geometry*. Birkhäuser Boston Inc., Boston, MA, 1997.
- [10] W. W. Crawley-Boevey. Maps between representations of zero-relation algebras. *J. Algebra*, 126(2):259–263, 1989.
- [11] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations ii: Applications to cluster algebras, 2009.
- [12] G. Dupont. Transverse quiver grassmannians and bases in affine cluster algebras, 2009. arXiv.org:0910.5494.
- [13] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529 (electronic), 2002.
- [14] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [15] Sergey Fomin and Andrei Zelevinsky. Cluster algebras: notes for the CDM-03 conference. In *Current developments in mathematics, 2003*, pages 1–34. Int. Press, Somerville, MA, 2003.
- [16] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. IV. Coefficients. *Compos. Math.*, 143(1):112–164, 2007.
- [17] David Hernandez and Bernard Leclerc. Cluster algebras and quantum affine algebras. arXiv.org:0903.1452.
- [18] Giovanni Cerulli Irelli. Canonically positive basis of cluster algebras of type $a_2^{(1)}$, 2009. arXiv.org:0904.2543.
- [19] Giovanni Cerulli Irelli. Quiver grassmannians associated with string modules, 2009. arXiv.org:0910.2592.
- [20] Bernhard Keller. Cluster algebras, quiver representations and triangulated categories, 2008. arXiv.org:0807.1960.
- [21] Leopold Kronecker. Algebraische reduction der schaaren bilinearer formen. *Sitzungsberichte Akad. Berlin*, pages 1225–1237, 1890.
- [22] Hiraku Nakajima. Quiver varieties and cluster algebras, 2009. arXiv.org:0905.0002.
- [23] Claus Michael Ringel. Exceptional modules are tree modules. *Proceedings of the Sixth Conference of the International Linear Algebra Society (Chemnitz)*.
- [24] Paul Sherman and Andrei Zelevinsky. Positivity and canonical bases in rank 2 cluster algebras of finite and affine types. *Mosc. Math. J.*, 4(4):947–974, 982, 2004.
- [25] Csaba Szanto. On the cardinalities of kronecker quiver grassmannians, 2009. arXiv.org:0903.1928.